

A prospective surveillance study of factor VIII inhibitor development in the Canadian haemophilia A population following the switch to a recombinant factor VIII product formulated with sucrose

M. RUBINGER,* D. LILICRAP,† G. E. RIVARD,‡ J. TEITEL,§ M. CARCAO,¶ C. HENSMAN,† I. WALKER** and THE ASSOCIATION OF HEMOPHILIA CLINIC DIRECTORS OF CANADA (AHCDC)¹

*CancerCare Manitoba, Department of Medical Oncology and Haematology, University of Manitoba, Winnipeg, MB;

†Department of Pathology and Molecular Medicine, Queen's University, Kingston; ‡Hôpital Sainte-Justine, Department of Pediatrics, Université de Montréal, Montréal, QC; §St Michael's Hospital, University of Toronto; ¶Division of

Hematology/Oncology, Department of Pediatrics, The Hospital for Sick Children, University of Toronto, Toronto; and

**McMaster University Medical Centre, Hamilton, ON, Canada

Summary. The introduction of new factor concentrates has, at times, resulted in an increase in inhibitor development; hence large systematic surveys of inhibitor development are necessary whenever new products are introduced. This study presents the results of a surveillance study conducted by the Inhibitor Subcommittee of the Association of Hemophilia Clinic Directors of Canada that evaluated inhibitor development in patients with haemophilia A following the switch to a second generation recombinant FVIII product (rFVIII-FS; Kogenate[®] Bayer). Four hundred and sixty haemophilia A paediatric and adults patients from 17 Canadian Comprehensive Hemophilia Care Centers were enrolled in the study. Of these, 274 patients had evaluable data. Blood samples collected at baseline (prior to the switch to rFVIII-FS), and at 12 and 24 months following conversion were tested for

inhibitors by the Nijmegen-modified Bethesda assay. Four subjects had positive inhibitor titres at baseline, with values ranging from 3.3 to 160 BU. Of the 274 patients who had baseline samples collected, 225 had postswitch samples collected at 12 months and 189 subjects had samples collected at 24 months. Only patients with positive baseline inhibitor titres ($n = 4$) had positive inhibitor titres at either the 12- or 24-month postswitch time points; therefore no *de novo* inhibitors developed over the 2-year evaluation period in this patient population. The results of this surveillance study suggest that the altered formulation of this recombinant FVIII concentrate was not associated with an increased incidence of inhibitor formation.

Keywords: factor VIII, haemophilia, inhibitor, surveillance

¹Association of Hemophilia Clinic Directors of Canada (AHCDC), Suite 2-008, 38 Shuter Street, Toronto, ON M5B 1A6, Canada. Tel.: +1 416 864 5042; fax: +1 416 864 5251; e-mail: ahcdc@smh.toronto.on.ca

Correspondence: Morel Rubinger, MD, CancerCare Manitoba, 675 McDermot Avenue, Winnipeg Room 2083, MB, R3E 0V9, Canada.

Tel.: +1 204 787 3594;

fax: +1 204 786 0196;

e-mail: morel.rubinger@cancercare.mb.ca

Accepted after revision 28 November 2007

Introduction

Haemophilia A is an inherited bleeding disorder caused by a deficiency of coagulation factor VIII (FVIII) that affects between 1/5000 to 1/10 000 males. The development of an inhibitor to FVIII (an antibody that neutralizes the coagulant activity of factor) following FVIII replacement therapy is the most serious treatment-related complication currently facing haemophilia patients; an inhibitor reduces the effectiveness of treatment, resulting in an increase in medical costs, and an increase in morbidity and mortality [1]. Known or suspected risk factors for the development of inhibitors to FVIII include: the

severity of disease, the genetic mutation responsible for haemophilia, family history of inhibitors, ethnicity, age of first exposure to FVIII, molecular modifications of the FVIII molecule and the number of exposure days to FVIII [2,3]. The incidence of inhibitors appears to vary among users of different FVIII concentrates, but there is no evidence to support the concern that switching from one product to another is itself a risk factor for inhibitor formation, independent of the FVIII product [4–8]. Recombinant FVIII products have the inherent safety benefit of eliminating the need for large pools of donor plasma, yet lingering concerns regarding the potential immunogenicity of recombinant products remain. That recombinant proteins can induce antibodies when given therapeutically is well illustrated by the occurrence of pure red cell aplasia induced by anti-erythropoietin antibodies following therapy with certain preparations of recombinant erythropoietin [9].

Formed in 1994, the Association of Hemophilia Clinic Directors of Canada (AHCDC) provides a structure through which Canadian haemophilia treaters, blood system regulators and operators can exchange information regarding product tracking, utilization, monitoring and surveillance for product efficacy and safety. Such monitoring is particularly important with the introduction of any new coagulation products. As reported in the study by Giles *et al.* [10], this organization initiated an inhibitor surveillance programme designed to address the theoretical concern that highly purified plasma-derived or recombinant FVIII products might be more immunogenic than earlier plasma derived products (this coincided with the conversion of most Canadian haemophilia A patients to either recombinant or affinity-purified plasma-derived preparations in 1994). An important element of the surveillance study design was the establishment of a central laboratory for the tracking and monitoring of inhibitors. The use of a central laboratory helped to ensure consistent methodology and standardized measurement for inhibitor detection, allowing evaluation and pooling of results across participating centres. In the study by Giles *et al.*, 478 patients switched from plasma-derived products to a first generation rFVIII product (Kogenate® Bayer, Bayer Healthcare, Berkeley, CA, USA) and inhibitor formation was then monitored for 1–2 years. This study found no evidence of increased inhibitor formation in these patients following the switch.

Similar to many other recombinant proteins, first generation rFVIII products, as studied by Giles *et al.* [10], were stabilized with human albumin in their final formulations. However, concerns regarding the

therapeutic use of mammalian-derived protein, such as human albumin, prompted the Medical and Scientific Advisory Council of the National Hemophilia Foundation in the US to encourage manufacturers to remove albumin from products used in the treatment of haemophilia [11]. Subsequently, Bayer Inc. developed a full-length rFVIII (Kogenate® FS; Bayer) that contains sucrose rather than albumin in the final formulation (rFVIII-FS) [12].

We report here a continuation of the efforts of the Inhibitor Subcommittee of the AHCDC, specifically evaluating inhibitor development following the conversion of haemophilia A patients to rFVIII-FS.

Materials and methods

Eligible subjects were Canadian paediatric and adults patients with moderate or severe haemophilia A who were switched from FVIII to rFVIII-FS. The study was approved by the respective review board/ethics committees of participating centres. This study was funded by Canadian Blood Services and Héma-Quebec following a recommendation from the AHCDC. Participation in the study was not influenced by the factor VIII product used prior to the switch, or concentrate history over the year prior to conversion. In addition, patients were eligible irrespective of whether an inhibitor was detected at baseline. The characteristics of the 274 eligible patients are summarized in Table 1. Based on FVIII measurements at baseline, 72.3% of patients were severe, 13.9% were moderate and 13.9% were mild. To be eligible for evaluation

Table 1. Patient characteristics*.

Age at switch to rFVIII-FS	
Mean age ± SD (years)	16.8 ± 10.2
Range (years)	0.9–40.8
Severity of haemophilia based on CRF data†	
Severe	220 (89)
Moderate	19 (8)
Mild	3 (1)
Severity not reported	4 (2)
Severity of haemophilia based on baseline FVIII measurement	
Severe (≤ 0.01 U mL ⁻¹)	198 (72)
Moderate (>0.01 – 0.05 U mL ⁻¹)	38 (14)
Mild (>0.05 U mL ⁻¹)	38 (14)
Family history of inhibitor†	
Yes	24 (10)
No	203 (83)
Unknown	19 (8)

Not all patients had completed CRFs; however, lab analyses were conducted on all samples collected, unless otherwise noted. Values are given as *n* (%).

**n* = 274 evaluable patients.

†*n* = 246 evaluable patients with completed CRF.

patients had to have baseline plasma samples collected within 3 months prior to the switch to rFVIII-FS and to have samples collected at 12 and 24 months following the switch to rFVIII-FS. All samples were drawn at least 48 h following any FVIII replacement therapy. Patients were withdrawn from the study if samples were not collected within 3 months of the 12- and 24-month postswitch time frame.

Blood samples were collected directly into vacuum-sealed tubes or indirectly via syringe and transferred into vacuum-sealed tubes. Platelet-poor plasma was obtained by centrifugation, and samples were frozen (-60°C or lower), and shipped to the Central Laboratory (Haemophilia Research Reference Laboratory, Kingston General Hospital, Kingston, ON) for analysis. All samples were tested for inhibitors by the Nijmegen-modification of the Bethesda method [13]. A positive inhibitor value was considered to be ≥ 0.5 BU.

Results

Four hundred and sixty haemophilia A patients from 17 Canadian comprehensive haemophilia care centres were enrolled. Of these, 274 met enrollment protocol requirements. During the time frame of this study, 28 August, 2000 until 28 September, 2003, an unanticipated disruption of rFVIII-FS production occurred (29 September 2001), and therefore some patients were switched from rFVIII-FS to other rFVIII products to manage the shortage. Data from such patients were included until the date they switched from rFVIII-FS to another product.

Study criteria were set out to include only moderate and severe haemophilia patients, but baseline factor measurements resulted in some patients being

recategorized as mild ($\text{FVIII} > 0.05 \text{ U mL}^{-1}$), in contrast to information on the case report forms (CRF) that categorized these patients differently. This discrepancy between baseline laboratory factor levels and the CRF may be explained by patients having received factor VIII 48–96 h prior to the baseline sample being taken or by simple imprecision of results from local laboratories; patients with levels of factor of $0.05\text{--}0.07 \text{ U mL}^{-1}$ may at times be found to have levels of 0.03 or 0.04 U mL^{-1} . Because the goal of this study was an evaluation of inhibitor development (a safety endpoint), mild, moderate and severe haemophilia patients were included in the data analysis. For most patients (82.5%) there was no family history of a FVIII inhibitor.

Subjects were excluded from the study for the following: problems with obtaining baseline sample (sample not obtained, $n = 7$, sample obtained after switch to rFVIII-FS, $n = 9$, sample obtained more than 90 days prior to switch to rFVIII-FS, $n = 47$), and problems with obtaining postswitch samples (samples not obtained, $n = 137$). As well, two subjects were excluded as they did not switch to rFVIII-FS ($n = 2$). As some patients had more than one exclusion criteria the final number of eligible patients amounted to 274 (Fig. 1).

While the goal of the study was to follow all patients for at least 2 years following conversion to rFVIII-FS, data were not collected for all patients at each of the protocol designated sampling times, both for reasons of non-availability of rFVIII-FS and other reasons. Of the 274 patients who had baseline samples collected, 12-month postswitch samples were collected for 225 and 24-month postconversion samples were collected for 189 subjects. One hundred and forty patients completed all three samples;




Fig. 1. Cohort of patients enrolled, excluded* and reason for exclusion, and patients eligible for analysis. *Patients have more than one exclusion criteria, which explains the discrepancy in the final number of patients eligible for analysis.

Table 2. Patients with positive inhibitor titres at baseline.

Patient	Baseline sample (BU)	12 months postswitch to rFVIII-FS (BU)	24 months postswitch to rFVIII-FS (BU)
1	3.3	1.2	NA
2	16	3.2	NA
3	160	33.8	27.7
4	4.5	4.6	5.8

BU, Bethesda units; NA, samples not collected for assay.

baseline, 12 and 24 months following conversion to rFVIII-FS. A slightly smaller group, 123 patients, completed all three samples, and used only rFVIII-FS replacement factor (without switching to another product) for the duration of the study period. Four subjects had positive inhibitor titres at baseline, with values ranging from 3.3 to 160 BU (Table 2). Of these, two patients had severe haemophilia, one had moderate haemophilia and for one patient the severity of haemophilia was unknown. Inhibitor assays for these four patients remained positive at 12 months following the switch to rFVIII-FS, with values ranging from 1.2 to 33.8 BU. At the 24-month postconversion time point, two of these patients tested positive for inhibitors (27.7 and 5.8 BU), while the remaining two subjects did not have 24-month samples collected. None of these patients received immune-tolerization.

Table 3 summarizes the inhibitor results from all valid patients for 12 and 24 months following conversion to rFVIII-FS. Only patients with positive baseline inhibitor titres ($n = 4$) had positive inhibitor titres at either the 12- or 24-month postswitch time points; therefore, no *de novo* inhibitors developed

over the 2-year evaluation period in this patient population. Specifically, in patients in whom all sequential samples were collected, there was no evidence of inhibitor development over the course of the study (Table 4). Similarly the 123 patients who received only rFVIII-FS during the study and had all the required samples collected did not show any evidence of inhibitor formation.

Discussion

The results of this surveillance study suggest that the formulation of recombinant FVIII with sucrose (Kogenate[®]-FS; Bayer) rather than albumin did not result in an increased risk of inhibitor formation in previously treated haemophilia A patients. The first surveillance study of the Canadian haemophilia A population showed that there was no increase in the incidence of FVIII inhibitors when previously treated patients (PTPs) were converted to either a first generation rFVIII or high purity affinity-purified plasma-derived FVIII [10].

It is important to emphasize that this surveillance study differed from a more structured clinical trial with regard to sampling frequency for inhibitor detection. Several clinical studies evaluating inhibitor formation in both previously untreated patients (PUPs) and PTPs collected samples at 3-month intervals for inhibitor titres [14–19]. In fact, in one study of PUPs and minimally treated patients, the frequency of sampling for inhibitor detection was even higher during the high-risk period [16]. With the less frequent sampling of every 12 months in this surveillance study, detection of transient inhibitors

Table 3. Inhibitor summary following conversion to rFVIII-FS*.

Baseline samples (preswitch)	12 months postswitch to rFVIII-FS			24 months postswitch to rFVIII-FS		
	Negative	Positive	Missing	Negative	Positive	Missing
Negative for inhibitors ($n = 270$)	221 (81%)	0	49 (18%)	185 (68%)	0	85 (31%)
Positive for inhibitors ($n = 4$)	0	4 (1.5%)	0	0	2 (0.7%)	2 (0.7%)†

*All samples, $n = 274$. A 'missing' sample can be a reflection of no sample collected at time point, or a patient switched to another product.

†Twenty-four month postswitch samples were collected for only two patients, with the other two being not evaluable due to 'missing' samples. A positive FVIII inhibitor had a value ≥ 0.5 BU.

Table 4. Inhibitor summary following conversion to rFVIII-FS – patients completing the full surveillance protocol*.

Baseline sample (preswitch)	12 months postswitch to rFVIII-FS			24 months postswitch to rFVIII-FS		
	Negative	Positive	Missing	Negative	Positive	Missing
Negative for inhibitors ($n = 138$)	138 (99%)	0	0	136 (97%)	0	2 (1.4%)
Positive for inhibitors ($n = 2$)	0	2 (1.4%)	0	0	2 (1.4%)	0

*Defined as subjects in whom all three samples (baseline, 12 and 24 months following conversion to rFVIII-FS) were obtained, $n = 140$. A 'missing' sample can be a reflection of no sample collected at specific time point, or a patient switched to another product. A positive FVIII inhibitor had a value ≥ 0.5 BU.

might have been missed. In addition, the study was not designed to match clinical evidence of an inhibitor (using FVIII recovery values or other clinical parameters) with laboratory detection, nor was it developed to detect non-neutralizing antibodies to FVIII. However, it is reasonable to suggest that the switch to rFVIII-FS from other recombinant FVIII formulations does not appear to lead to the development of new inhibitors of important clinical significance.

The relationship between FVIII product type and inhibitor risk is clearly an important issue for haemophilia patients and care givers, and the subject of ongoing debate [4–6]. An association between modification of the FVIII production (plasma-derived, pasteurized FVIII concentrates with either prior controlled-pore silica adsorption or solvent detergent treatment) and an increased incidence of inhibitors in PTPs was documented by Peerlinck *et al.* and Rosendaal *et al.* [20,21]. The pasteurization process may have produced epitope alterations in these preparations that resulted in the increased development of inhibitors. A recent study in France compared inhibitor incidence in PUPs treated with either a single recombinant or a single plasma-derived FVIII product [22]. These investigators noted a lower incidence of inhibitors associated with the use of the plasma-derived FVIII product compared with rFVIII. However, review of several studies with either plasma-derived or rFVIII products suggests that the recombinant products are not more immunogenic than FVIII preparations when the comparisons take into consideration the details of study design (including the frequency of inhibitor testing) and risk factors influencing inhibitor development [7,8]. *In toto* these data suggest that individual FVIII molecules may possess different inhibitor-inducing profiles but that amongst the many risk factors known to affect inhibitor development any one factor may be difficult to isolate. Also relevant to a discussion of inhibitor incidence is the number of exposure days (EDs), as patients with <20 EDs are still at high risk for inhibitor formation [7]. While the number of EDs was not documented in the present surveillance study, most of the patients enrolled in this study had received many more than 20 EDs. Additionally, it is important to note the recommendation of the Scientific Subcommittee of the International Society of Thrombosis and Haemostasis to use PTPs as the appropriate population in which to evaluate product immunogenicity [23].

Differences in study design, numerous risk factors for inhibitor development (severity of haemophilia, genetic mutation type, ethnicity, number of EDs,

etc.) and the heterogeneity of the patient population complicate direct comparison of inhibitor incidence between studies. While it is difficult to prospectively assess one specific host or treatment-related risk factor, it is important to continually monitor new and existing FVIII replacement products for inhibitor development, and to identify significant deviations from the very low frequency 'background' immunogenicity of these products.

Acknowledgements

In addition to the author, the following Canadian Hemophilia Clinic Directors participated: Dr Dorothy Barnard (Halifax); Dr Victor Blanchette (Toronto); Dr Mason Bond (Vancouver); Dr Josée Brossard (Sherbrooke); Dr Manuel Caracao (Toronto); Dr Robert Card (Saskatoon); Dr Anthony K.C. Chan (Hamilton); Dr Stephanie Cloutier (Quebec); Dr Jeffrey Davis (Vancouver); Dr Michael Delorme (Kelowna); Dr Christine Demers (Quebec); Dr Sean Dolan (St John's); Dr Nancy Dower (Edmonton); Dr Bernadette Garvey (Toronto); Dr Kulwant Gill (Sudbury); Dr Jack Hand (St John's); Dr Rosemary Henderson (Charlottetown); Dr Donald S. Houston (Winnipeg); Dr Sara Israels (Winnipeg); Dr Lawrence Jardine (London); Dr Paula James (Kingston); Dr Robert Klassen (Ottawa); Dr Mariette Lepine-Martin (Sherbrooke); Dr David Lillicrap (Kingston); Dr Reinhard Lohmann (London); Dr Patricia J. McCusker (Winnipeg); Dr Man-Chiu Poon (Calgary); Dr Victoria Price (Halifax); Dr Margaret L. Rand (Toronto); Dr Bruce Ritchie (Edmonton); Dr Georges Rivard (Montreal); Dr Sue Robinson (Halifax); Dr Sheldon Rubin (Moncton); Dr Mary France Scully (St John's); Dr Sam Schulman (Hamilton); Dr Mariana Silva (Kingston); Dr Jean St-Louis (Montreal); Dr Jerome Teitel (Toronto); Dr Alan Timmouh (Ottawa); Dr Dimitrios Vergidis (Thunder Bay); Dr Linda Vickars (Vancouver); Dr Irwin Walker (Hamilton); Dr Margaret Warner (Montreal); Dr Phillip Wells (Ottawa); Dr Blair Whitemore (Montreal); Dr Rochelle Winikoff (Montreal); Dr John W.Y. Wu (Calgary); Dr John K. M. Wu (Vancouver); Dr Margaret Yhap (Halifax). I wish to thank Wendy Scales, PhD, Scientific Norton, LLL, for editorial contribution and Pamela Johnston, RN, for her diligence in data collection.

Disclosures

Dr. Teitel has acted as a paid consultant for Baxter Healthcare, Wyeth and Novo Nordisk. He is a

recipient of research grant support from the Bayer Partnership fund. All other authors stated that they had no interests which might be perceived as posing a conflict or bias.

References

- 1 Key NS. Inhibitors in congenital coagulation disorders. *Br J Haematol* 2004; **127**: 379–91.
- 2 Lusher J. Natural History of inhibitor development in children with severe hemophilia A treated with factor VIII products. In: Lee Blackwell C, Berntorp E, Hoots W, eds. *Textbook of Hemophilia*. Malden, Massachusetts: Publishing Ltd., 2005: 34–9.
- 3 Dimichele D, Rivard G, Hay C, Antunes S. Inhibitors in haemophilia: clinical aspects. *Haemophilia* 2004; **10**(Suppl. 4): 140–5.
- 4 Aledort LM. Is the incidence and prevalence of inhibitors greater with recombinant products? Yes. *J Thromb Haemost* 2004; **2**: 861–2.
- 5 Hoots WK, Lusher J. High-titer inhibitor development in hemophilia A: lack of product specificity. *J Thromb Haemost* 2004; **2**: 358–9.
- 6 Lusher JM. Is the incidence and prevalence of inhibitors greater with recombinant products? No. *J Thromb Haemost* 2004; **2**: 863–5.
- 7 Scharrer I, Bray GL, Neutzling O. Incidence of inhibitors in haemophilia A patients – a review of recent studies of recombinant and plasma-derived factor VIII concentrates. *Haemophilia* 1999; **5**: 145–54.
- 8 Scharrer I, Ehrlich HJ. Lack of evidence for increased inhibitor incidence in patients switched from plasma-derived to recombinant factor VIII. *Haemophilia* 2001; **7**: 346–8.
- 9 Boven K, Stryker S, Knight J *et al.* The increased incidence of pure red cell aplasia with an Eprex formulation in uncoated rubber stopper syringes. *Kidney Int* 2005; **67**: 2346–53.
- 10 Giles AR, Rivard GE, Teitel J, Walker I. Surveillance for factor VIII inhibitor development in the Canadian Hemophilia A population following the widespread introduction of recombinant factor VIII replacement therapy. *Transfus Sci* 1998; **19**: 139–48.
- 11 Medical Scientific and Advisory Council (MASAC). *Recommendation Regarding the Use of Recombinant Factor VIII in the Treatment of Hemophilia A*. New York: National Hemophilia Foundation, 1995, Medical Bulletin 232 (Chapter Advisory # 234).
- 12 Boedeker BG. Production processes of licensed recombinant factor VIII preparations. *Semin Thromb Hemost* 2001; **27**: 385–94.
- 13 Verbruggen B, Novakova I, Wessels H, Boezeman J, van den Berg M, Mauser-Bunschoten E. The Nijmegen modification of the Bethesda assay for factor VIII:C inhibitors: improved specificity and reliability. *Thromb Haemost* 1995; **73**: 247–51.
- 14 Abshire T, Brackmann H-H, Scharrer I *et al.* Sucrose-formulated recombinant human antihemophilic factor VIII is safe and efficacious for treatment of hemophilia A in home therapy: results of a multicenter, international, clinical investigation. *Thromb Haemost* 2000; **83**: 811–6.
- 15 Bray GL, Gomperts ED, Courter S *et al.* A multicenter study of recombinant factor VIII (Recombinate): safety, efficacy, and inhibitor risk in previously untreated patients with hemophilia A. *Blood* 1994; **83**: 2428–35.
- 16 Kreuz W, Gill JC, Rothschild C *et al.* Full-length sucrose-formulated recombinant factor VIII for treatment of previously untreated or minimally treated young children with severe haemophilia A. Results of an international clinical investigation. *Thromb Haemost* 2005; **93**: 457–67.
- 17 Lusher JM, Lee CA, Kessler CM, Bedrosian CL. The safety and efficacy of B-domain deleted recombinant factor VIII concentrate in patients with severe haemophilia A. *Haemophilia* 2003; **9**: 38–49.
- 18 Lusher J, Abildgaard C, Arkin S *et al.* Human recombinant DNA-derived antihemophilic factor in the treatment of previously untreated patients with hemophilia A: final report on a hallmark clinical investigation. *J Thromb Haemost* 2004; **2**: 574–83.
- 19 White GC II, Courter S, Bray GL *et al.* A multicenter study of recombinant factor VIII (Recombinate) in previously treated patients with hemophilia A. The Recombinate Previously Treated Patient Study Group. *Thromb Haemost* 1997; **77**: 660–7.
- 20 Peerlinck K, Arnout J, DiGiambattista M, Gilles J, Laub R, Jacquemin M. Factor VIII inhibitors in previously treated haemophilia A patients with a double virus-inactivated plasma derived Factor VIII concentrate. *Thromb Haemost* 1997; **77**: 80–6.
- 21 Rosendaal FR, Nieuwenhuis HK, van den Berg HM *et al.* A sudden increase in factor VIII inhibitor development in multitransfused hemophilia A patients in The Netherlands. Dutch Hemophilia Study Group. *Blood* 1993; **81**: 2180–6.
- 22 Goudemand J, Rothschild C, Demiguel V *et al.* Influence of the type of factor VIII concentrate on the incidence of factor VIII inhibitors in previously untreated patients with severe hemophilia A. *Blood* 2006; **107**: 46–51.
- 23 White GC, DiMichele DM, Mertens K *et al.* Utilization of previously treated patients (PTPs), noninfected patients (NIPs), and previously untreated patients (PUPs) in the evaluation of new Factor VIII and Factor IX concentrates. *Thromb Haemost* 1999; **81**: 462.