



# Pediatric Bleeding Questionnaire (PBQ) Scoring Key

Date:

Patient name:

Patient number:

Score Symptom	-1	0	1	2	3	4
<b>Epistaxis</b>	-	No or trivial ( $\leq 5$ per year)	$>5$ per year OR $>10$ minutes duration	Consultation only	Packing, cauterization or antifibrinolytics	Blood transfusion, replacement therapy or desmopressin
<b>Cutaneous</b>	-	No or trivial ( $\leq 1$ cm)	$>1$ cm AND no trauma	Consultation only	-	-
<b>Minor wounds</b>	-	No or trivial ( $\leq 5$ per year)	$>5$ per year OR $>5$ minutes duration	Consultation only or Steri-strips	Surgical hemostasis or antifibrinolytics	Blood transfusion, replacement therapy or desmopressin
<b>Oral cavity</b>	-	No	Reported at least once	Consultation only	Surgical hemostasis or antifibrinolytics	Blood transfusion, replacement therapy or desmopressin
<b>Gastrointestinal tract</b>	-	No	Identified cause	Consultation or spontaneous	Surgical hemostasis, antifibrinolytics, blood transfusion, replacement therapy or desmopressin	-
<b>Tooth extraction</b>	No bleeding in at least 2 extractions	None done or no bleeding in 1 extraction	Reported, no consultation	Consultation only	Resuturing, repacking or antifibrinolytics	Blood transfusion, replacement therapy or desmopressin
<b>Surgery</b>	No bleeding in at least 2 surgeries	None done or no bleeding in 1	Reported, no consultation	Consultation only	Surgical hemostasis or antifibrinolytics	Blood transfusion, replacement therapy or desmopressin
<b>Menorrhagia</b>	-	No	Reported or consultation only	Antifibrinolytics or contraceptive pill use	D&C or iron therapy	Blood transfusion, replacement therapy, desmopressin or hysterectomy
<b>Post-partum</b>	No bleeding in at least 2 deliveries	No deliveries or no bleeding in 1 delivery	Reported or consultation only	D&C, iron therapy or antifibrinolytics	Blood transfusion, replacement therapy or desmopressin	-
<b>Muscle hematoma</b>	-	Never	Post-trauma, no therapy	Spontaneous, no therapy	Spontaneous or traumatic, requiring replacement therapy or desmopressin	Spontaneous or traumatic, requiring surgical intervention or blood transfusion
<b>Hemarthrosis</b>	-	Never	Post-trauma, no therapy	Spontaneous, no therapy	Spontaneous or traumatic, requiring replacement therapy or desmopressin	Spontaneous or traumatic, requiring surgical intervention or blood transfusion
<b>Central nervous system</b>	-	Never	-	-	Subdural, any intervention	Intracerebral, any intervention
<b>Other</b> Post-circumcision Umbilical stump Cephalohematoma Macroscopic hematuria Post-venepuncture Conjunctival hemorrhage	-	No	Reported	Consultation only	Surgical hemostasis, antifibrinolytics or iron therapy	Blood transfusion, replacement therapy or desmopressin

Total Bleeding Score \_\_\_\_\_